

November 2005

Universality and Quark Masses of the Staggered Fermion Action

Anna Hasenfratz*

Department of Physics, University of Colorado, Boulder, CO-80304-0390

Staggered fermions with 4 tastes are expected to describe 4-flavor QCD in the continuum limit, therefore at finite lattice spacing the staggered determinant should be equivalent to an $SU(4)$ flavor-symmetric system up to lattice artifacts. This equivalence is the starting point of the 4th root trick used to reduce the number of fermion flavors and provides the only consistent description of 2 or 1 flavor systems. In this paper we argue that the quark mass of the underlying flavor symmetric theory differs from the staggered mass by an additive term due to the taste breaking of the staggered action. The relation is the same for 2 and 1 flavor fermions. This additive mass shift implies that at finite lattice spacing staggered simulations correspond to heavier quark masses than indicated by the staggered Goldstone pion and that staggered fermions cannot reach the chiral limit at finite lattice spacing.

I. NOTE ADDED

The justification of the 4th root trick is essential for the large scale high precision simulations carried out in recent years using 2+1 flavors staggered sea quarks. The validity of the procedure depends on the properties of the staggered action. In this paper I assumed that the unrooted 4-taste staggered action differs from a flavor degenerate 4-flavor action only in local pure gauge terms and investigated the consequences of this assumption.

*Electronic address: anna@eotvos.colorado.edu;

Since the original publication of this manuscript there is growing evidence indicating that this assumption is incorrect [1, 2][17]. In a recent paper the author and Hoffmann [2] presented numerical evidence in the 2 dimensional Schwinger model that indicates that while the rooted staggered action might not be local at finite lattice spacing, in the continuum limit the non-local terms become irrelevant and the rooted action can describe chiral fermions with arbitrary physical mass. The same paper also justifies the ad-hoc assumption of this manuscript about the mass shift between the physically matched staggered and overlap fermions. I would like to emphasize that this mass offset is a lattice artifact. For further details the reader is referred to the publication [2] and references there.

II. INTRODUCTION

Staggered fermion lattice actions have four non-degenerate flavors or tastes. There is little doubt that in the continuum limit they correspond to QCD with four degenerate flavors. Consider a renormalization group transformation that respects gauge and chiral symmetry, with a fixed point on the $m = 0$ and $g^2 = 0$ critical surface that represents 4-flavor continuum QCD. Along the renormalized trajectory of the RG transformation the lattice actions describe the same continuum system; these actions presumably have Ginsparg-Wilson chiral, but at least $SU(4)$ flavor symmetry. The renormalization group transformation should move the staggered lattice action toward this renormalized trajectory, and since RG transformations change the action only in irrelevant, cut-off level terms, the $SU(4)$ flavor symmetric action along the renormalized trajectory will differ from the original staggered one only in irrelevant, cut-off level terms.

In this paper we investigate this underlying $SU(4)$ flavor symmetric action at finite lattice spacing. We argue that its bare fermion mass must differ from the staggered mass not only in a multiplicative but in an additive mass renormalization factor as well. This additive term is related to the taste breaking of the staggered action and vanishes only in the continuum limit. An immediate consequence of this observation is that the staggered

Goldstone pion can be significantly lighter than the flavor symmetric pions, leading to incorrect mass estimates at finite lattice spacing. We support our argument with data from the 2-dimensional Schwinger model.

III. DECOMPOSITION OF THE STAGGERED DETERMINANT

The 4-taste staggered determinant describes 4-flavor QCD in the continuum limit if its determinant can be decomposed as

$$\det(D_{\text{st}}) = \det(D_{\text{s4}})\det(T) , \quad (1)$$

where D_{st} is the staggered Dirac operator, D_{s4} is an $\text{SU}(4)$ flavor symmetric lattice operator, and the operator T gives cut-off level corrections only. If one could rigorously show that the action $\tilde{S} = -\text{Tr} \ln(T)$ was a local operator in the continuum limit, Eq.1 would validate the 4th root procedure used in reducing the number of flavors in staggered simulations [3, 4, 5, 6, 7]. We are not concerned about this issue right now, so all we assume is that T does not change the physical predictions.

Both D_{st} and D_{s4} has one relevant operator corresponding to the fermion mass. (The relevant direction in the gauge field is governed by the gauge action which is not specified in Eq.1.) We will denote the corresponding bare masses by m_q and m_{s4} . The renormalized quark masses differ from the bare ones by multiplicative renormalization factors.

Eq.1 comes from a RG based operator relation, it has to be valid on all, or at least most, gauge configurations. Consider a topologically non-trivial configuration at finite lattice spacing. In the chiral $m_{\text{s4}} = 0$ limit D_{s4} has 4 degenerate zero modes, $\det(D_{\text{s4}}) = 0$. On the other hand D_{st} has no exact zero modes at any finite lattice spacing, $\det(D_{\text{st}}) \neq 0$ even when $m_q = 0$. The determinant of the cut-off level term T has to be finite, so Eq.1 cannot be valid if both $m_q = 0$ and $m_{\text{s4}} = 0$. Yet Eq.1 is necessary if 4-taste staggered fermions are in the same universality class as 4-flavor continuum QCD. The only way to satisfy the

equation is to allow m_{s4} to remain finite when the staggered quark mass vanishes, i.e.

$$m_{s4} = a^2 \Delta + Z m_q. \quad (2)$$

The additive correction is $O(a^2)$ but present at any finite lattice spacing.

Eq.1 suggests two different ways to look at 4-taste staggered fermions. The first is the usual staggered interpretation: the action describes four fermionic flavors but due to lattice artifacts the flavor symmetry is broken, only a remnant $U(1)$ symmetry survives. At the massless $m_q = 0$ limit the system has one Goldstone pion, the other pseudoscalars have $O(a^2)$ masses. Only in the continuum limit do we recover the full $SU(4) \times SU(4)$ flavor symmetry. The second interpretation is based on the right hand side of Eq.1. It states that the theory has 4 degenerate fermions, it differs from a Ginsparg-Wilson action only in cut-off terms, so in the $m_{s4} = 0$ limit we expect 15 massless pions. The two interpretations are compatible but their corresponding fermionic fields are very different. They have different lattice artifacts and their predictions can also differ by $O(a^2)$ effects, but in the continuum limit they are identical. For a consistent field theory one has to use the same action for the valence as for the sea quarks. For 4-taste staggered fermions we could use either the staggered action or the $SU(4)$ flavor symmetric action in the valence sector.

The relation between the staggered and flavor symmetric quark masses is the same for 1-flavor staggered fermions defined by the 4th root of the staggered determinant, assuming the 4th root does not introduce non-universal, non-local terms. There is an important difference though between the 4-taste and 1-flavor cases. The 4th root procedure is correct only if we consider the $SU(4)$ flavor symmetric interpretation, so for a consistent field theoretical description one has to use this 4th root operator in the valence quark sector as well. When the standard staggered operator is used in the valence sector we have to deal with partial-quenching/mixed actions effects, and also a mismatch between the staggered and flavor symmetric quark masses. Problems due to partial-quenching/mixed actions clearly show up when one attempts to calculate the pseudoscalar singlet [8, 9]. One should note that these inconsistencies are not related to the 4th root determinant being non-local,

they simply show the inconsistencies between the sea and valence quark actions.

Staggered chiral perturbation theory [10, 11, 12] models the 4th root by hand correcting the number of fermions in the sea quark loops. That approach assumes that the 4th root of the staggered Dirac operator is a local theory. We know that this is not the case [13]. For a consistent perturbative treatment one should consider staggered valence quarks on the background of Ginsparg-Wilson quarks whose mass is adjusted/fitted appropriately. Mixed action simulations has to match the valence and sea quark masses. Again, this matching should be done to the $SU(4)$ flavor symmetric masses.

A finite additive mass implies that with staggered fermions one cannot reach the chiral limit at any finite lattice spacing. This might explain the contradictions of the finite temperature scaling behavior of $N_f = 2$ simulations. For these systems the $O(4)$ critical point is beyond the reach of the simulations, it lies at negative quark mass, though it likely influences the scaling behavior at positive mass values as well. At the same time there is an $O(2)$ invariant critical point at $m_q = 0$ that mainly effects the fermionic observables. The combination of these two nearby critical points can make the scaling of the system very complicated.

IV. NUMERICAL SUPPORT

In a series of papers Durr and Hoelbling investigated the 2-dimensional Schwinger model with 1 and 2 flavors both with staggered and overlap fermions [5, 14]. Here we consider their results for the topological susceptibility.

In Ref.[14] the topological susceptibility was measured through the index of the massless overlap operator, so any deviation between the staggered and overlap action is due to the difference between the dynamical configurations. Chiral perturbation theory predicts that the topological susceptibility vanishes in the chiral limit. The solid curves in Fig. 1 show the susceptibility both on $N_f = 1$ and 2 overlap configurations. Both curves follow the expected behavior. The diamonds and octagons on the left panel show the result




Figure 1: The topological susceptibility in the 2D Schwinger model with $N_f = 1$ and 2 flavors with overlap and staggered dynamical fermions. The left panel is from Ref.[14]. The right panel is the same data replotted with the quark mass for the staggered fermions shifted as $m = m_q + 0.065$.

on staggered dynamical configurations [18]. The susceptibility remains finite even at the $m_q \rightarrow 0$ limit indicating that the underlying chiral theory has a finite mass. Re-plotting the staggered data with a mass shift $m_q \rightarrow m_q + 0.065$ (right panel of Fig. 1) brings the overlap and staggered curves together. The agreement is not perfect but very good. Including a multiplicative Z factor like in Eq.2 as $m = 1.05m_q + 0.055$ makes the agreement excellent everywhere but at the lowest $m_q = 0$ point. The fact that both the $N_f = 1$ and 2 data set requires the same mass shift indicates that both are governed by the same chiral action with $m \approx m_q + 0.065$. This observation supports the expectation that the square root action of $N_f = 1$ is a correct 1-flavor action. The data in Fig. 1 corresponds to unsmeared fermions. Even one level of APE smearing reduces the difference between the staggered and overlap actions, the required mass shift becomes $m - m_q \lesssim 0.01$. Of course the taste violation is also similarly reduced after a smearing step.

There were other unexpected differences between the staggered and overlap results in Refs. [5, 14], mainly with the scalar condensate. It is likely that those too will be resolved by observing the mass shift between the two actions.

One would like to estimate the mass shift for the 4D Asqtad action. Measurements of the topological susceptibility clearly indicate a mass shift [15]. Staggered chiral perturbation theory predicts that the topological susceptibility scales with the taste-singlet pion[16], suggesting that the additive mass is governed by the heaviest of the staggered pions.

V. CONCLUSION

We argued that if 4-taste staggered fermions describe 4-flavor QCD in the continuum limit, at any finite lattice spacing the fermion mass of the underlying chiral 4-flavor theory differs from the mass of the staggered action not only by a multiplicative but by an additive term as well. The same relation holds between the masses for 1 and 2-flavor systems.

This additive mass is a lattice artifact and will go away in the continuum limit. At finite lattice spacing however staggered measurements can underestimate the light hadron masses if those are extracted from the lowest mass staggered operators. The mass shift can have strong effects on finite temperature simulations and can explain the large lattice artifacts observed in the topological susceptibility as well. The additive mass should be taken into account in any simulation using mixed actions.

VI. ACKNOWLEDGMENTS

This work was supported by the US Department of Energy. I am indebted to T. DeGrand for numerous discussions. I would like to thank S. Sharpe and S. Durr for extensive

correspondence, P. Hasenfratz and F. Niedermayer for helpful questions and comments.

-
- [1] C. Bernard, M. Golterman, and Y. Shamir (2006), in preparation.
 - [2] A. Hasenfratz and R. Hoffmann (2006), hep-lat/0604010.
 - [3] D. H. Adams (2004), hep-lat/0411030.
 - [4] S. Dürer, C. Hoelbling, and U. Wenger, Phys. Rev. **D70**, 094502 (2004), hep-lat/0406027.
 - [5] S. Dürer and C. Hoelbling, Phys. Rev. **D71**, 054501 (2005), hep-lat/0411022.
 - [6] Y. Shamir, Phys. Rev. **D71**, 034509 (2005), hep-lat/0412014.
 - [7] C. Bernard et al. (2005), hep-lat/0509176.
 - [8] E. B. Gregory, A. C. Irving, C. McNeile, S. Miller, and Z. Sroczynski (2005), hep-lat/0509193.
 - [9] S. Prelovsek (2005), hep-lat/0510080.
 - [10] W.-J. Lee and S. R. Sharpe, Phys. Rev. **D60**, 114503 (1999), hep-lat/9905023.
 - [11] C. Aubin and C. Bernard, Phys. Rev. **D68**, 034014 (2003), hep-lat/0304014.
 - [12] C. Aubin and C. Bernard, Phys. Rev. **D68**, 074011 (2003), hep-lat/0306026.
 - [13] B. Bunk, M. Della Morte, K. Jansen, and F. Knechtli, Nucl. Phys. **B697**, 343 (2004), hep-lat/0403022.
 - [14] S. Dürer and C. Hoelbling, Phys. Rev. **D69**, 034503 (2004), hep-lat/0311002.
 - [15] C. Bernard et al., Phys. Rev. **D68**, 114501 (2003), hep-lat/0308019.
 - [16] B. Billeter, C. DeTar, and J. Osborn, Phys. Rev. **D70**, 077502 (2004), hep-lat/0406032.
 - [17] Many of these recent developments were discussed in the “Workshop on the 4th root of the staggered determinant” at INT, Seattle, March 2006.
 - [18] The author is indebted to S. Durr for providing the data for the plot.